

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Tilmicosol

*(in Cyprus, Czech Republic, Greece, Hungary, Spain, Lithuania, Portugal and United Kingdom)*

Tilmic

*(in Denmark)*

TylmiScan

*(in Poland)*

250 mg/ml Concentrate for Oral Solution for Use in Drinking Water or Milk Replacer

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

#### **Active substance:**

Tilmicosin (as phosphate) 250 mg/ml

#### **Excipients:**

Propyl gallate (E 310) 0.2 mg/ml

Disodium edetate 2.0 mg/ml

For the full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Concentrate for oral solution for use in drinking water or milk replacer.

Clear yellow to amber coloured solution.

### 4. CLINICAL PARTICULARS

#### **4.1 Target species**

Chickens (except hens producing eggs for human consumption),

Turkeys,

Pigs,

Calves (non ruminant)

## 4.2 Indications for use, specifying the target species

Pigs: For the treatment and metaphylaxis of respiratory disease in pig herds, associated with *Mycoplasma hyopneumoniae*, *Pasteurella multocida*, *Actinobacillus pleuropneumoniae*.

Chickens: For the treatment and metaphylaxis of respiratory disease in chicken flocks, associated with *Mycoplasma gallisepticum* and *M. synoviae*.

Turkeys: For the treatment and metaphylaxis of respiratory disease in turkey flocks, associated with *Mycoplasma gallisepticum* and *M. synoviae*.

Calves: For the treatment of bovine respiratory disease, associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Mycoplasma bovis*, *M. dispar*.

## 4.3 Contraindications

Do not allow horses and other equines access to drinking water containing tilmicosin.

Do not use in case of hypersensitivity to tilmicosin or to any of the excipients.

Do not administer to ruminating animals with active rumen function.

## 4.4 Special warnings for each target species

Important: Must be diluted before administration to animals

Pigs, chickens and turkeys: Water consumption should be monitored in order to guarantee adequate dosing. In case water consumption does not match quantities for which recommended concentrations were calculated, the concentration of Tilmicosol has to be adapted in a way that the recommend dosage will be taken up by the animals or different medication should be considered.

## 4.5 Special precautions for use

### Special precautions for use in animals

For oral use only. Contains disodium edetate; do not inject.

Severely ill individuals tend to drink less and may need simultaneous treatment, preferably by parenteral medication.

Inappropriate use of the product may increase the prevalence of bacteria resistant to tilmicosin and may decrease the effectiveness of treatment with tilmicosin-related substances. The use of the product should be based on susceptibility tests.

Medicated drinking water or milk replacer should be prepared fresh every 24 hours.

Official, national and regional antimicrobial policies should be taken into account when the product is used.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals

Tilmicosin may induce irritation. Macrolides, such as tilmicosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or

eye. Hypersensitivity to tilmicosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

To avoid exposure during preparation of the medicated drinking water, wear overalls, safety glasses, and impervious gloves. Do not eat, drink or smoke when handling this product. Wash hands after use.

In the case of accidental ingestion, wash out mouth immediately with water and seek medical advice. In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water. Do not handle the product if you are allergic to ingredients in the product.

If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

#### **4.6 Adverse reactions (frequency and seriousness)**

In very rare cases (less than 1 animal in 10,000 animals, including isolated reports), a decrease in water intake has been observed.

#### **4.7 Use during pregnancy, lactation or lay**

The safety of the veterinary medicinal product has not been established during pregnancy, lactation or lay. Use only accordingly to the benefit/risk assessment by the responsible veterinarian.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

None known.

#### **4.9 Amounts to be administered and administration route**

Pigs: To be included in the drinking water to provide a daily dose of 15-20 mg tilmicosin per kg bodyweight for 5 days, which may be achieved by the inclusion of 200 mg tilmicosin per litre (80 ml Tilmicosol per 100 litres).

Chickens and turkeys (except hens producing eggs for human consumption): To be included in the drinking water at a daily dose of 15-20 mg/kg bodyweight in chickens and 10-27 mg/kg bodyweight in turkeys for 3 days, which may be achieved by the inclusion of 75 mg tilmicosin per litre (30 ml Tilmicosol per 100 litres).

Calves: To be included in milk replacer only, at a dose of 12.5 mg/kg bodyweight and given twice daily for 3-5 consecutive days, which may be achieved by the inclusion of 1 ml of product every 20 kg bodyweight

Medicated drinking water should be prepared fresh every 24 hours using only clean water.

Medicated milk replacer should be prepared fresh every 8 hours using only clean water.

If signs of disease do not significantly improve within 3-5 days, the diagnosis should be re-evaluated and treatment changed.

To avoid underdosing body weight should be determined as accurately as possible. The uptake of medicated water depends on the clinical condition of the animals. In order to obtain the correct dosage the concentration of the product has to be adjusted accordingly.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

When pigs are offered drinking water containing 300 or 400 mg tilmicosin/litre (equivalent to 22.5-40 mg/kg bodyweight or 1.5-2 times the recommended concentration) commonly animals exhibit a reduced water intake. Although this has a self-limiting effect on tilmicosin intake, it could, in extreme circumstances, result in dehydration. This can be corrected by the removal of the medicated drinking water and replacement with fresh unmedicated water.

No symptoms of overdose have been seen in chickens given drinking water containing levels of tilmicosin up to 375 mg/litre (equivalent to 75-100 mg/kg bodyweight or 5 times the recommended dose) for 5 days, daily treatment with 75 mg/litre (equivalent to the maximum recommended dose) for 10 days resulted in a decrease in faecal consistency.

No symptoms of overdose have been seen in turkeys given drinking water containing levels of tilmicosin up to 375 mg/litre (equivalent to 50-135 mg/kg bodyweight or 5 times the recommended dose) for 3 days, daily treatment with 75 mg/litre (equivalent to the maximum recommended dose) for 6 days also produced no symptoms of overdose.

No symptoms of overdose, with exception of a slight decrease in the milk consumption, have been seen in calves given twice daily doses 5 times the maximum recommended dose or for twice the maximum recommended duration of treatment.

#### **4.11 Withdrawal period(s)**

Meat and offal of:

Pigs - 14 days

Chickens - 12 days

Turkeys - 19 days

Calves - 42 days

Not authorised for use in laying birds producing eggs for human consumption. Do not use within 14 days of onset of the laying.

Not authorised for use in lactating animals producing milk for human consumption.

## **5. PHARMACOLOGICAL PROPERTIES**

Pharmacotherapeutic Group: Antibacterials for systemic use, macrolides

**ATCVet code:** QJ01FA91

## 5.1 Pharmacodynamic properties

Tilmicosin is a semi-synthetic antibiotic of the macrolide group, and is believed to affect protein synthesis. It has bacteriostatic action but at high concentrations it may be bactericidal.

This antibacterial activity is predominantly against Gram-positive microorganism with activity against certain gram-negative ones and Mycoplasma of a bovine, porcine, ovine and avian origin. In particular, its activity has been demonstrated against the following microorganism:

- Pigs: *Mycoplasma hyopneumoniae*, *Pasteurella multocida* and *Actinobacillus pleuropneumoniae*
- Chickens and turkeys: *Mycoplasma gallisepticum* and *Mycoplasma synoviae*
- Calves: *Mannheimia haemolytica*, *Pasteurella multocida*, *Mycoplasma bovis*, *M. dispar*

Scientific evidence suggests that macrolides act synergistically with the host immune system. Macrolides appear to enhance phagocyte killing of bacteria. Tilmicosin has been shown to inhibit *in vitro* the replication of the Porcine Reproductive and Respiratory Syndrome virus in alveolar macrophages in a dose dependent fashion.

Cross-resistance between tilmicosin and other macrolides and lincomycin has been observed.

## 5.2 Pharmacokinetic particulars

Whilst blood concentrations of tilmicosin are low, there is pH-dependent macrophage accumulation of tilmicosin in inflamed tissue.

Pigs: After oral administration of 200 mg tilmicosin/litre drinking water, the average active substance concentrations detected in lung tissue, alveolar macrophages and bronchial epithelium 5 days after the start of treatment were reported 1.44 µg/ml, 3.8 µg/ml and 7.4 µg/g respectively.

Poultry: As early as 6 hours after oral administration of 75 mg tilmicosin/litre drinking water, the average active substance concentrations detected in lung and alveolar tissue were 0.63 µg/g and 0.30 µg/g respectively. 48 hours after the start of the treatment, the tilmicosin concentrations in lung and alveolar tissue were 2.3 µg/g and 3.29 µg/g respectively.

Calves: As early as 6 hours after oral administration of 25 mg tilmicosin/kg bodyweight/day in milk replacer, an average active substance concentration of 3.1 µg/g was detected in lung tissue. 78 hours after the start of treatment, the tilmicosin concentration in lung tissue

was 42.7 µg/g. Therapeutically effective concentrations of tilmicosin were measured up to 60 hours after treatment.

Turkeys: After oral administration of 75 mg tilmicosin/litre drinking water, the average active substance concentrations detected in lung tissue, air sac tissue and plasma 5 days after the start of treatment were 1.89 µg/ml, 3.71 µg/ml and 0.02 µg/g respectively. The highest mean tilmicosin concentration detected for lung tissues was 2.19 µg/g at 6 days; for air sac tissue it was 4.18 µg/g at 2 days and in the plasma it was 0.172 µg/g at 3 days.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Disodium edetate

Propyl gallate, E 310

Phosphoric acid, concentrated (for pH adjustment)

Purified water

### **6.2 Incompatibilities**

None known.

### **6.3 Shelf life**

Shelf-life of the veterinary medicinal product as packaged for sale: 2 years

Shelf-life after first opening the immediate packaging: 3 months

Shelf-life after dilution or reconstitution according to directions: 24 hours

### **6.4. Special precautions for storage**

Do not store above 25°C. Store in the original container in order to protect from light. Do not freeze.

### **6.5 Nature and composition of immediate packaging**

250 ml in white high density polyethylene bottle with polypropylene screw cap

1000 ml in white high density polyethylene bottle with polypropylene screw cap and calibrated polypropylene measuring cup

Not all pack sizes may be marketed.

### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Manure from treated animals should not be deposited on the same field in successive years.

**7. MARKETING AUTHORISATION HOLDER**

Lavet Pharmaceuticals Ltd.

1161 Budapest,

Ottó u. 14.

Hungary

**8. MARKETING AUTHORISATION NUMBER**

Vm 32823/4003

**9. DATE OF FIRST AUTHORISATION**

11 February 2011

**10. DATE OF REVISION OF THE TEXT**

February 2016